


EUROPEAN JOURNAL OF  
**MOLECULAR MEDICINE**



## **European Journal of Molecular medicine**

**Volume 2, No.5, September 2022**

**Internet address:** [tp://ejournals.id/index.php/EJMM/issue/archive](http://ejournals.id/index.php/EJMM/issue/archive)

**E-mail:** [info@ejournals.id](mailto:info@ejournals.id)

Published by ejournals PVT LTD

DOI prefix: 10.52325

Issued Bimonthly

Potsdamer Straße 170, 10784 Berlin, Germany

### *Requirements for the authors.*

*The manuscript authors must provide reliable results of the work done, as well as an objective judgment on the significance of the study. The data underlying the work should be presented accurately, without errors. The work should contain enough details and bibliographic references for possible reproduction. False or knowingly erroneous statements are perceived as unethical behavior and unacceptable.*

*Authors should make sure that the original work is submitted and, if other authors' works or claims are used, provide appropriate bibliographic references or citations. Plagiarism can exist in many forms - from representing someone else's work as copyright to copying or paraphrasing significant parts of another's work without attribution, as well as claiming one's rights to the results of another's research. Plagiarism in all forms constitutes unethical acts and is unacceptable. Responsibility for plagiarism is entirely on the shoulders of the authors.*

*Significant errors in published works. If the author detects significant errors or inaccuracies in the publication, the author must inform the editor of the journal or the publisher about this and interact with them in order to remove the publication as soon as possible or correct errors. If the editor or publisher has received information from a third party that the publication contains significant errors, the author must withdraw the work or correct the errors as soon as possible.*

### **OPEN ACCESS**

Copyright © 2022 by Thematics Journals of Applied Sciences

## CHIEF EDITOR

**Serikuly Zhandos PhD,**

Associate Professor, RWTH Aachen University, Aachen, Germany

## EDITORIAL BOARD

**Bob Anderson**  
ImmusanT, *USA*

**Marco Bruno**  
Erasmus Medical Center,  
*The Netherlands*

**Antoni Castells**  
Hospital Clinic  
*Barcelona, Spain*

**Giacomo Caio**  
University of Ferrara, *Italy*

**Michael Farthing**  
St George's Hospital Medical  
School, *UK*

**Carmelo Scarpignato**  
University of Parma,  
*Italy*

### Geriatric Medicine

**Ian Cameron**  
The University of Sydney,  
*Australia*

**Sutthichai Jitapunkul**  
Chulalongkorn University,  
*Thailand*

**Juulia Jylhävä**  
Karolinska Institute, *Sweden*

**Kenneth Rockwood**  
Dalhousie University,  
*Canada*

ELSEVIER



SSRN  
European Research Portal

Universal  
Impact Factor



**SURGICAL TREATMENT OF SPINE HEMANGIOMAS****Rakhmatov K.R.**Bukhara State Medical Institute,  
Uzbekistan, Bukhara city  
dr\_karim\_neyroxirurg@mail.ru

*Abstract: The results of surgical treatment of 27 patients with aggressive hemangiomas of the thoracic and lumbar spine were analyzed. All patients were conditionally divided into two groups, taking into account the severity of the pathological process. Vertebroplasty with bone cement was common in the treatment of both groups of patients. Differences in the groups were in one group, taking into account the presence of an intracanal component, compression of the dural sac and nerve roots, it was the performance of decompression-stabilizing transpedicular osteosynthesis of the spine with open vertebroplasty. Obtaining almost identical ratios of good and satisfactory results of treatment in both groups of patients confirms that the algorithm of surgical treatment used and the amount of treatment performed in one or two operations, taking into account the severity of the pathological process, were reasonable and correct.*

*Keywords: Hemangioma, vertebroplasty, neurological, deficit, pain syndrome, hemangioma, local pain, radicular syndromes, health.*

**ХИРУРГИЧЕСКОЕ ЛЕЧЕНИЕ ГЕАНГИОМ ПОЗВОНОЧНИКА.****Рахматов К.Р.**Бухарской Медицинский Институт,  
Узбекистан, г. Бухара.  
dr\_karim\_neyroxirurg@mail.ru

*Аннотация: Проанализированы результаты хирургического лечения 27 больных с агрессивными гемангиомами грудного и поясничного отдела позвоночника. Все больные условно разделены с учетом выраженности патологического процесса на две группы. Общим в лечении обеих групп больных являлось выполнение вертебропластики костным цементом. Отличия в группах заключались в одной группе, учитывая наличия внутриканального компонента, компрессию дурального мешка и нервных корешков являлось выполнение декомпрессионно-стабилизирующего транспедикулярного остеосинтеза позвоночника с открытой вертебропластикой. Получение практически одинаковых соотношений хороших и удовлетворительных результатов лечения в обеих группах больных подтверждает, что используемый алгоритм хирургического лечения и выполнение объема лечения за одну или две операции с учетом выраженности патологического процесса были обоснованными и правильными.*

*Ключевые слова: гемангиома, вертебропластика, неврологический, дефицит, болевой синдром, гемангиома, локальная боль, корешковые синдромы, здоровье.*

Introduction. Until now, vertebral hemangioma is still an incompletely studied nosological form, neither the pathogenesis of the disease has been determined, nor the optimal treatment tactics have been developed. And today, in most hospitals, radiation therapy remains the main method of treating aggressive hemangiomas, the effectiveness of which is extremely low. Hemangiomas of the vertebral bodies, detected at autopsies

in 5%-11% of patients, are the most common benign neoplasms in which vertebroplasty is performed (1,5,7). Currently, the following types of treatment for spinal hemangiomas are used: surgery, transarterial embolization, percutaneous vertebroplasty, radiation therapy (8,9,10). During surgical treatment, there is a high risk of massive intraoperative bleeding and non-radical removal of the formation. A long recovery period is also one of the unfavorable factors when choosing a treatment method. Transarterial embolization causes partial occlusion of the supplying arterial vessels of the hemangioma, but does not cause destruction of the formation itself. Partial entry of the sclerosing material into the intercostal or lumbar arteries can cause pain in these patients. If necessary, re-embolization is not possible due to occlusion of the feeding vessels. Radiation therapy is a fairly effective method of treatment, however, the effect of radiation is delayed in time, and the risk of post-radiation myelopathy is also high. (2,3,4,6).

The aim of the study was to improve the quality of life of patients with vertebral hemangiomas using open and through skin vertebroplasty.

Materials and methods. The work is based on the analysis of 27 cases of patients with aggressive hemangiomas of the spine, who were examined and treated at the Bukhara Regional Multidisciplinary Medical Center.2018 to 2020

Of the studied 27 patients, there were 12 (44.4%) men and 15 (55.6%) women. The average age of the patients was 45 and ranged from 28 to 62 years. The distribution of patients by gender and age is presented in Table 1.

Table 1

## The distribution of patients by gender and age

Age in years	Gender				Total	
	Male		Female			
	abs.	%	abs.	%	abs.	%
<b>21-44 years old</b>	3	11	6	22,2	9	33,2
<b>45-59 years old</b>	7	26	5	18,6	12	44,6
<b>Age 60 and above</b>	2	7,4	4	14,8	6	22,2
<b>Total</b>	12	44,44	15	55,6	27	100

The table above shows that female patients predominated in all age groups. The largest age group consisted of patients 45-59 years old 12 (44.6) and 21-44 years old 9 (33.2%), that is, the most able-bodied age.

We assessed the intensity of the pain syndrome using a four-point DENIS scale (1989), where the criteria were the use of painkillers. The distribution of patients depending on the intensity of the pain syndrome is presented in Table 2.

Table 2

**Distribution of patients depending on the intensity of pain  
according to the DENIS scale**

Points*	Number of patients	%
0	0	0
1	8	29,6
2	14	51,8
3	5	18,6
4	0	0
Total	27	100

\* Note: 0 - no pain; 1 - minimal or rare pain that does not require anesthesia; 2 - moderate pain relieved by non-narcotic analgesics; 3 - pain of moderate intensity, in which it is periodically necessary to prescribe narcotic analgesics; 4 - unbearable pain, requiring regular prescription of narcotic analgesics.

In most cases, the pain was constant, periodically increased or decreased for no apparent reason, sometimes it was associated with vertical physical activity. Very often, patients were disturbed by night pains. Physiotherapy, which was often prescribed to such patients, aggravated the pain.

The condition of all patients, both at admission and at discharge and in the long term, was assessed using the Karnovsky quality of life scale. According to the Karnovsky scale, the quality of life of patients was assessed in the long-term period after surgery based on the obtained follow-up data.

According to the scale, 100-80% is regarded as a satisfactory condition, 70-40% - moderate, and 30-10% - a serious condition. Upon admission, 21 (77.8%) patients were in a satisfactory condition, 6 were in a moderate condition ( 22.2%) patients in serious condition were not observed.

Interventions were performed at Th3-L5 levels. Percutaneous vertebroplasty was performed in the thoracic region - 6 vertebrae, in the lumbar region - 10 vertebrae, in the thoracic and lumbar regions - 12 vertebrae. Operational aid - decompressive laminectomy with posterior fusion with transpedicular fixation systems and open vertebroplasty was performed in 6 (22.2%) patients. Needles were passed through transpeduncular access. The puncture of the vertebral bodies was performed under image intensifier tube (IIT) control. The technique of vertebroplasty was no different from that generally accepted today. Bone cement was injected using Stryker PCD systems with the addition of contrast agents (barium sulfate powder).

In all cases, control spondylography was performed.






Fig. 1. Hemangioma of the VL1 vertebral body: a - spondylogram of the lumbar spine in lateral projection; b - MRI of the lumbar spine; c - MSCT after decompression-stabilizing surgery and open vertebroplasty.

Surgical treatment was in 6 (22.2%) patients. Decompressive-stabilizing operations with transpedicular fixation with plates with open vertebroplasty: in the thoracic region - 4 (14.8%) and lumbar - 2 (7.4%) patients.

Results and discussion. Aggressive vertebral hemangiomas were clinically manifested in most cases by persistent pain syndrome of moderate intensity (2 points according to DENIS) without significant differences in the compared groups.

Local pain syndrome, reflex muscular-tonic reactions, limited movements in the spine and general motor activity of patients during the day. However, in most cases (6 cases), motor activity disorders did not reach the degree at which the patient would not be able to move without assistance. In this connection, the scale for assessing the quality of life did not reflect the entire range of options for changes in the motor activity of patients with vertebral hemangiomas. In no case (in the second group) did the score of motor activity disorders exceed one according to the J.R. Gaughen. That is, there were no patients in the sample who were in a forced horizontal position or moved with the help of a wheelchair. The frequency of admission and the choice of the pharmacological group of painkillers directly depended on the severity of local pain and radicular

syndromes and on individual characteristics of pain tolerance. The rest periodically or constantly used non-steroidal anti-inflammatory drugs.

Of all 27 (100%) operated patients, 22 (81.4%) patients noted the absence of pain, 5 (18.6%) decreased pain. In the postoperative period, 85.1% of operated patients noted a significant regression of pain.

Improvement of the neurological status after surgical treatment in the second group occurred in 6 (100%) patients, the neurological deficit improved slightly. The Karnofsky scale was used to assess the functional outcome of treatment. At the same time, good results suggested a decrease in neurological deficit and an increase in the Karnofsky index in the early postoperative period.

Analysis of the immediate results of surgical treatment, on average 10-14 days after surgery, shows that out of all 27 (100%) patients, good results on the Karnofsky scale were achieved in 25 (92.6%), satisfactory results in 2 (7.4 %) observations, unsatisfactory results, i.e. there was no persistent deterioration in the condition of patients. Assessment of immediate results on the Karnofsky scale indicates a positive impact of surgery on the health and well-being of patients in most cases.

#### Conclusions.

1. Indications for open vertebroplasty with decompression-stabilizing surgery in this category of patients were: intense pain syndrome, the presence of an intracanal hemangioma component, compression of the dural sac, neurological deficit.

2. Puncture vertebroplasty is a more effective treatment for aggressive spinal hemangiomas. This method helps to reduce the intensity of the pain syndrome.

3. Puncture vertebroplasty is a minimally invasive technique with a minimum percentage of complications, which significantly reduces the length of the patient's stay in the hospital. After vertebroplasty puncture, the increase in strength properties of vertebral bodies and the normalization of venous hemodynamics contribute not only to the regression of pain, but also to an increase in movement activity in the spine and resistance to physical stress in the first week after the surgery.

#### Used literature.

1. Balberkin A.V., Morozov A.K., Shavyrin D.A. Long-term results of surgical treatment of benign tumors of the thoracic and lumbar spine in adults. *Vestn. injuries. and an orthopedist. N.N. Priorov.* - 2004. - No. 2. - S. 63-67.

2. Voronovich I.R., Pashkevich L.A. *Spinal tumors.* - Minsk, 2000. - 240 p.

3. Zharkov P.L., Zharkov A.P., Bubnovsky S.M. *Lumbar pain. Diagnostics. The reasons. Treatment.* - M.: Medicine, 2000. - 143 p.

4. Slyn'ko E.I. *Surgical treatment of vascular tumors of the spine and spinal cord // Ukrainian neurosurgical journal.* - 2000. - No. 1 (9). - S. 55-64.

5. Asumu T.O., Williamson B., Hughes D.G. *Symptomatic spinal hemangiomas in association with cutaneous hemangiomas. A case report // Spine.* - 1996. - 21. - P. 1082-1084.

6. Bavbek M., Yurt A., Caner H., Altinors N. *An unusual dumbbell form of cavernous hemangioma of the cervical spine // Kobe J. Med. Sci.* - 1997. - 43 (2). - P. 57-63.

7. Dahlin, D.C. *Bone tumors: General Aspects and Data on 6221 cases*, 3rd edition / D.C. Dahlin. - Springfield, IL: Charles C. Thomas, 1978.

8. Galibert, P. *Preliminary note on the treatment of vertebral angioma by percutaneous acrylic vertebroplasty / P. Galibert, H. Deramond*

9. Fox M.W., Onofrio B.M. *The natural history and management of symptomatic and asymptomatic vertebral hemangiomas // J. Neurosurg.* - 1993. - 78. - P. 36-45.