


EUROPEAN JOURNAL OF
MOLECULAR MEDICINE



European Journal of Molecular medicine

Volume 3, No.5, October 2023

Internet address: <http://ejournals.id/index.php/EJMM/issue/archive>

E-mail: info@ejournals.id

Published by ejournals PVT LTD

DOI prefix: 10.52325

Issued Bimonthly

Potsdamer Straße 170, 10784 Berlin, Germany

Requirements for the authors.

The manuscript authors must provide reliable results of the work done, as well as an objective judgment on the significance of the study. The data underlying the work should be presented accurately, without errors. The work should contain enough details and bibliographic references for possible reproduction. False or knowingly erroneous statements are perceived as unethical behavior and unacceptable.

Authors should make sure that the original work is submitted and, if other authors' works or claims are used, provide appropriate bibliographic references or citations. Plagiarism can exist in many forms - from representing someone else's work as copyright to copying or paraphrasing significant parts of another's work without attribution, as well as claiming one's rights to the results of another's research. Plagiarism in all forms constitutes unethical acts and is unacceptable. Responsibility for plagiarism is entirely on the shoulders of the authors.

Significant errors in published works. If the author detects significant errors or inaccuracies in the publication, the author must inform the editor of the journal or the publisher about this and interact with them in order to remove the publication as soon as possible or correct errors. If the editor or publisher has received information from a third party that the publication contains significant errors, the author must withdraw the work or correct the errors as soon as possible.

OPEN ACCESS

Copyright © 2023 by Thematics Journals of Applied Sciences

CHIEF EDITOR

Serikuly Zhandos PhD,

Associate Professor, RWTH Aachen University, Aachen, Germany

EDITORIAL BOARD

Bob Anderson

ImmusanT, *USA*

Marco Bruno

Erasmus Medical Center,
The Netherlands

Antoni Castells

Hospital Clinic
Barcelona, Spain

Giacomo Caio

University of Ferrara, *Italy*

Michael Farthing

St George's Hospital Medical
School, *UK*

Carmelo Scarpignato

University of Parma,
Italy

Geriatric Medicine

Ian Cameron

The University of Sydney,
Australia

Sutthichai Jitapunkul

Chulalongkorn University,
Thailand

Juulia Jylhävä

Karolinska Institute, *Sweden*

Kenneth Rockwood

Dalhousie University,
Canada

**MORPHOLOGY OF LYMPHOPOIETIC FUNCTION OF REGIONAL LYMPH NODES OF BACTERIAL INFECTION IN PURULENT PYELONEPHRITIS
RSSPMCU, RPPC**

**J.A. Shadmanov
R. Israilov**

Abstract: In this scientific research, morphological changes characteristic of the lymphocytopoietic function of regional lymph nodes in cystitis and pyelonephritis were studied. If in the initial period of the disease the barrier and drainage function of lymphatic vessels and regional lymph was increased, over time it is observed that B-lymphocytic lymphocytopoietic function prevails in regional lymph nodes. In the morphological aspect, the lymphocytopoietic function is confirmed by activation and increase by proliferation of lymphoid and macrophage cells in the cortical layer of lymph nodes.

Keywords: pyelonephritis, cystitis, drainage, lymphocytopoiesis, lymph vessel, lymph node, sinusoid, morphofunctional zone, proliferation.

The relevance of the topic. Pyelonephritis in 80% of cases is caused by E. coli, this bacillus normally lives in the intestine, enters the urinary tract in violation of the hygiene of the pelvis and calyx, as well as the interstitial tissues [8]. As a result, starting from the renal tissue, it causes stagnation in the tissue of venous and lymphatic vessels. When the development of this process begins, due to the development of compensatory processes, pyelo-renal reflux occurs, the calyx and pelvis expand. As a result, lymphangitis and lymphadenitis develop from the spread of infection to the lymphatic capillaries of the kidneys and ureters. Lymphadenitis can acquire a lymphoproliferative character in response to antigens of microorganisms. As humoral immunity to bacterial infection develops, lymphocytes proliferate in the lymph node, mainly in the B-lymphocytic zone, and lymphocytopoiesis develops [6].

Lymphocytopoiesis, which is the process of formation, proliferation and differentiation of lymphoid cells during the formation of lymphocytes, is the development of all lymphoid cells synchronously with each other [1, 2]. Lymphocytopoiesis proceeds in stages both in the embryonic and postpartum periods. Lymphocytopoiesis, in turn, consists of T-lymphocytopoiesis and B-lymphocytopoiesis. Each of them consists of three stages: the bone marrow stage, which occurs in antigen-independent central organs, and antigen-dependent stages, which occur in peripheral lymphoid organs.

In B-lymphocytopoiesis, B-lymphocytes first mature in the bone marrow, then migrate to the spleen and lymph nodes, differentiate in them and accumulate in a zone specific to B-lymphocytes, that is, in lymphoid follicles, germinal, perilymphoid ring and marginal areas. B-lymphocytes detects antigens with the participation of a macrophage and a T helper, identifies the antigen and synthesizes immunoglobulin in response to the antigen [3, 4]. For the synthesis of immunoglobulin, the B-lymphocyte turns into a plasma cell, and this process is called "B-lymphocyte plasmation". In this process, macrophages, monocytes, histiocytes and a dendritic cell phagocytose, process and determine antigens that have entered the tissue, and help lymphocytes detect them using histomolecular antigens on their outer surface.

Immature T-lymphocytes in T-lymphocytopoiesis are formed in the bone marrow, then migrate to the cortical layer of the thymus, where, in a sterile state, they turn into cortical thymocytes on average within a week. After a week, 2-4% of cortical thymocytes

survive, the rest (96-98%) undergo apoptosis and phagocytosis by thymus macrophages. Macrophages determine the products of phagocytized thymocytes, distribute them to the surviving thymocytes, and they differentiate. These surviving thymocytes should be able to identify their own histomolecular antigen on the surface of other cells and perform a state of innate immunological tolerance.

The selection of surviving thymocytes with such high requirements prevents their transition to a state of autoaggression in relation to private cells in the body, in violation of which autoimmune diseases develop [5, 7]. When antigen enters the body, it is determined by macrophages and is presented to the T-lymphocyte, T-lymphocytes undergo a "blastotransformation" reaction, turning into an activated lymphocyte, that is, an immunocyte, and exhibit cytotoxic action against the antigen, destroying it.

The purpose of the study. This scientific research was aimed at studying morphological changes characteristic of lymphocytopoietic function in regional lymph nodes in patients with purulent pyelonephritis.

Material and methods. To achieve this goal in 2021-2022 during the autopsy of corpses of patients with purulent pyelonephritis, lymph nodes located in the pelvis were taken for histological examination in the department of adult Pathology at the Center of Republican Pathological Anatomy of the Ministry of Health of the Republic of Uzbekistan. The lymph nodes were divided into two equal parts and hardened within 72 hours in 10% neutralized formalin dissolved in a phosphate buffer. Then it was washed in running water for 3-4 hours, dehydrated in a series of alcohols with an increased concentration and poured paraffin with wax, prepared blocks. Histological sections of 5-6 microns were made from paraffin blocks, stained with hematoxylin-eosin dyes, examined in the lenses of a light microscope 20, 40, studied, and took computer pictures of the desired areas.





Research results and their discussion. Examination of the lymph nodes of the inguinal region at the microscopic level in chronic purulent cystitis and pyelonephritis showed that, since this disease is a bacterial infectious disease, lymphocytopoietic function prevails in the lymph nodes, that is, lymphoproliferative inflammation. Macroscopically, it was found that the lymph nodes of the inguinal region are enlarged to varying degrees, that is, the smallest is 854 mg, and the largest is 1760 mg.

It was found that the lymph nodes have a bean-shaped and rounded appearance, the outer surface is smooth, but the color has changed due to inflammation, i.e. it has acquired a brownish hue due to the presence of red, bluish and grayish foci. When cut, the texture is soft, with a rotten appearance, the color is spotty.

In histological preparations treated with hematoxylin and eosin dyes by the general morphological method, in the initial period of the disease with purulent pyelonephritis, there is a sharp swelling of the interstitial substance in all morphofunctional areas of the lymph node, expansion of the subcapsular sinusoid, loosening of the tissue structures of the fibrous membrane, rarefaction of cells. It is claimed that the predominance of dyscirculatory and edematous processes coincides with the exudative period of inflammatory disease.

Primary follicles located in the mucous membrane of the lymph node are loosened and are defined as enlarged in area, with a sparse arrangement of lymphocytes, macrophages and reticular cells. The presence in the composition of lymphocytes of cells with different sizes of the nucleus and dark and light color indicates increased lymphocytopoietic activity in them (Fig. 1). It is known that from the location of B-lymphocytes in the lymphoid follicles of the mucous membrane of the lymph nodes, it is revealed that the plasmatic reaction of B-lymphocytes develops to develop humoral immunity in response to bacterial infection.

In the initial periods of the disease with purulent pyelonephritis, there is a sharp expansion of the sinusoids of both the cortical and cerebral layers, the presence of fragments of tissues and cells, microorganisms in the cavity due to the predominance of the processes of dyscirculation and edema in the lymph node. As the lymphocytopoietic reaction begins to develop intensively in the parenchyma of the lymph node, proliferation of lymphoid and monocyte-macrophage cells is detected both in the lymphoid follicles of the cortex and in the paracortical layer (Fig. 2).

	
<p>Figure 1. Primary lymphoid follicle of the cortical layer of the lymph node, the interstitial substance of which is strongly swollen, lymphocytes are located in a loosened and hyperchromasized form. Staining: G-E. Increased: 10x40.</p>	<p>Figure 2. General view of the lymph node tissue, at the initial stage of purulent pyelonephritis, the sinusoids are dilated, the cortex and paracortical layer are hyperplasiized. Staining: G-E. Increased: 10x10.</p>
	
<p>Figure 3. View of the lymph node in a semi-thin section, in a state of activation of large and medium lymphocytes. Staining: toluidine blue. Increased: 10x100.</p>	<p>Figure 4. Proliferation and increase of B-lymphocytes in the lymph node of the inguinal region in chronic purulent pyelonephritis. Staining: G-E. Increased: 10x10.</p>

When semi-thin sections 300-400 nm long were made for a deeper study of the lymphopoietic function of lymphoid cells in the lymph node, it was found that the reticular cells that make up the stroma of the lymph node tissue were relatively hypertrophied and enlarged in size. In particular, it is revealed that the nuclei of reticular cells are enlarged in size, have an oblong and oval shape, the karyoplasm is colored pale, the nucleus is hypertrophied, fine concentrations are formed in the heterochromatin karyolemma. Among the reticular cells, the location of lymphocytes is

ELSEVIER
 SSRN
 Universal Impact Factor

observed, which are activated and differentiated to varying degrees.

The fact that most of these lymphocytes consist of large and medium-sized cells was judged by the structure of their nucleus (Fig. 3). There was a sharp hypertrophy of the nucleolus of large lymphocytes, characteristic of the activation of their nuclei. It was found that heterochromatin forms a uniform concentrated row at the edge of the nucleus adjacent to the karyolemma. It is observed that medium-sized lymphocytes turn into immunoblasts with increased proliferation and differentiation. Their activation was manifested by hyperchromasia of the nucleus, hypertrophy of the nucleus, and an increase in heterochromatin.

In chronic purulent pyelonephritis, lymphatic vessels and lymph nodes first develop barrier and drainage function, and then the predominance of lymphocytopoietic function in the lymph nodes is detected. Morphologically, hyperplasia and an increase in the size of the lymphoid follicles of the cortical layer of the lymph nodes are observed. In lymphoid follicles, an increase in the lymphocytopoietic reaction is observed, manifested in an increase in cell size due to cell hyperplasia in the germinal center of proliferation with an increase in the surrounding zones of the B-lymphocytic ring (Fig.4).

The fact that the germinal center is mostly round in shape and of different sizes indicates hypertrophy of secondary lymphoid follicles.

In bacterial infectious purulent pyelonephritis, antigen-dependent lymphocytopoiesis develops in response to bacteria in the lymph nodes, in which there is an appearance, a shift to the side, the formation and differentiation of lymphocytes in the germinal center.

Morphologically, lymphoid follicles turn into a secondary form, in the center of which there is a germinative region in which dendritic reticular cells are activated, causing the proliferation of lymphocytes, contributing to their improvement. The picture shows a secondary lymphoid follicle, in which there are hypertrophied large-shaped dendritic reticular cells, among which the number of lymphoblasts and large lymphocytes is increased. The lymphocytes differentiated and formed a lymphocytic ring around the germinative region (Fig. 5), in which the lymphocytes are arranged in rows.




Figure 5. Secondary follicle of the inguinal lymph node in chronic purulent pyelonephritis, an increase in the number of reticulocytes, lymphoblasts in the germinal center. Staining: G-E. Increased: 10x40.

Figure 6. Expansion of all morphofunctional zones of secondary lymphoid follicles. Staining: G-E. Increased: 10x40.

Over time, the zone of secondary lymphoid follicles expands and increases, in the germinal region it is observed that reticular cells with dendrites have acquired the appearance of a "milky way" due to the fact that they have turned into cells of a whitish

color. The morphofunctional state of the lymphoid follicles in this form indicates an increased lymphocytopoietic function in them. It turns out that the germinative zone has merged with the surrounding lymphocytic ring, and its zone has also expanded dramatically. Small lymphocytes in the lymphocytic ring are fully mature, formed lymphocytes that, as a result of migration to the medullar layer, undergo an immunological response, the so-called "plasmatisation" inside the sinusoid of the medullar layer and in the soft bands between them, turning into plasma cells and synthesizing immunoglobulins, forming antibodies corresponding to the antigens of existing microorganisms, and humoral immune response.

Conclusion

While at the initial stage of the development of bacterial infectious purulent pyelonephritis, the barrier and drainage function in the lymphatic vessels and regional lymph nodes increases, over time there is a predominance of the development of B-lymphocytic lymphocytopoietic function in the lymph nodes.

It was determined that in the initial period of the disease with purulent pyelonephritis, the processes of dyscirculation and edema prevail in the lymph node, the sinusoids of both the cortex and the cerebral layer are sharply expanded, fragments of tissues and cells, microorganisms are present in the cavity, and later in the parenchyma of the lymph node, a lymphocytopoietic reaction begins to develop intensively, both in the lymphoid follicles of the cortical layer and in the paracortical layer the proliferation of lymphoid and monocyte-macrophage cells begins.

At the beginning of the lymphocytopoietic reaction, the primary follicles first loosen and expand in area, the lymphocytes, macrophages and reticular cells contained in them are activated and sparsely located, hyperplasia of the lymph node mucosa in response to bacterial infection indicates the predominance of B-lymphocytopoiesis.

Over time, the zone of secondary lymphoid follicles expands and increases; the increased lymphocytopoietic function in it is evidenced by the observation that in the germinative zone, reticular cells with dendrites acquired the appearance of a "stellate sky" due to transformation into whitish cells.

Small lymphocytes in the lymphocytic layer are fully developed, formed lymphocytes that, as a result of migration to the medullar layer, undergo an immunological response, the so-called "plasmatisation" inside the sinusoid of the medullar layer and in the soft bands between them, turning into plasma cells and synthesizing immunoglobulins, forming antibodies corresponding to the antigens of existing microorganisms, and humoral immune response.

Reference

1. Alekseev N.A. Clinical aspects of leukopenia, neutropenia and functional disorders of neutrophils. St. Petersburg: Foliant, 2002: 416.
2. Bobova L.P., Kuznetsov S.L., Saprykin V.P. Histophysiology of blood and organs of hematopoiesis and immunogenesis. M.: New Wave, 2003: 157.
3. Gorchakov V.N., Gorchakova O.V., Gorchakov V.N. Functional morphology of lymph nodes of different localization and their response to phytotherapy // Modern problems of science and education. - 2015. - № 4. ;
4. Rumyantsev A. G. Syndrome of enlarged lymph nodes as a pediatric problem / A. G. Rumyantsev, V. M. Chernov, V. M. Delyagin // Attend. Doctor. - 1998. - No. 3. - pp. 49-52.
5. Ryjova O.B., Torubarova N.A. The role of viral infections in the pathogenesis of cytopenic syndromes in newborn children. Materials of the XI Congress "Man and medicine". M., 2004: 137-138.

6.The structure of the incidence of pyelonephritis / S.N. Styajkina et al.//Problems of modern science and education. - 2006, p.245.

7.Tsinslerling A.V., Tsinslerling V.A. Modern infections. Pathological anatomy and pathogenesis issues. 2nd ed. St. Petersburg: Sotis, 2002: 352.

8.Etiology and pathogenesis of pyelonephritis. Rumyantsev A.Sh., Goncharova N.S. Nephrology, 2015.pp. 40