



# TJAS

**Thematic Journal of Applied Sciences**

informing scientific practices around the world  
through research and development

# Thematic Journal of Applied Sciences

Volume 1, No 2, May 2021

Internet address: <http://ejournals.id/index.php/TJAS/issue/archive>

E-mail: [info@ejournals.id](mailto:info@ejournals.id)

Published by Thematics Journals PVT LTD

Issued Bimonthly

Chief editor

**S. G. Ahmed**

*Professor of Computational Mathematics and Numerical Analysis Faculty of Engineering, Zagazig University, Zagazig, Egypt, P. O. Box 44519*

*Requirements for the authors.*

*The manuscript authors must provide reliable results of the work done, as well as an objective judgment on the significance of the study. The data underlying the work should be presented accurately, without errors. The work should contain enough details and bibliographic references for possible reproduction. False or knowingly erroneous statements are perceived as unethical behavior and unacceptable.*

*Authors should make sure that the original work is submitted and, if other authors' works or claims are used, provide appropriate bibliographic references or citations. Plagiarism can exist in many forms - from representing someone else's work as copyright to copying or paraphrasing significant parts of another's work without attribution, as well as claiming one's rights to the results of another's research. Plagiarism in all forms constitutes unethical acts and is unacceptable. Responsibility for plagiarism is entirely on the shoulders of the authors.*

*Significant errors in published works. If the author detects significant errors or inaccuracies in the publication, the author must inform the editor of the journal or the publisher about this and interact with them in order to remove the publication as soon as possible or correct errors. If the editor or publisher has received information from a third party that the publication contains significant errors, the author must withdraw the work or correct the errors as soon as possible.*

**OPEN ACCESS**

Copyright © 2021 by Thematics Journals of Applied Sciences

## CHIEF EDITOR

**S. G. Ahmed**

*Professor of Computational Mathematics and Numerical Analysis Faculty of Engineering, Zagazig University, Zagazig, Egypt, P. O. Box 44519*

## EDITORIAL BOARD

**Yu Li**

*Wuhan University of Technology, China*

**Seung Man Yu**

*Seoul National University of Science and Technology, South Korea*

**Seyed Saeid Rahimian Koloor**

*Universiti Teknologi Malaysia, Malaysia*

**Eko Susanto**

*Menegment of journal Indonesia*

**Siti Mazlina Mustapa Kamal**

*Universiti Putra Malaysia, Malaysia*



## INSECTS AND THE INFLUENCE OF ENVIRONMENTAL FACTORS ON THEM.

**Khonkhodzhaeva Nodira Bakhtiyarovna**

Associate Professor, Tashkent State Pedagogical University named after Nizami,  
Republic of Uzbekistan. Tashkent city, Uzbekistan.  
E-mail: nodiraxanxadjueva@gmail.com

*Abstract: This article discusses the prospects for the use of pheromones in agriculture as a method of pest control and their advantages over pesticides from an environmental point of view. In this article, pheromones, their specific significance, as well as their significance for biological species, ways of their possible use for human needs, as well as the influence of external physical environmental factors on pheromones. In modern environmental conditions, this issue is most acute, and research in this area is very promising. Pheromones are one of the types of external stimuli that affect the behavior and physiological state of humans and animals, a complex of special olfactory signals is shown. These are biological markers of a kind, volatile signals that control neuroendocrine behavioral responses, developmental processes, and processes associated with social behavior and reproduction. Pheromones help alter the behavior, physiological and emotional state or metabolism of other individuals of the same species. Pheromones have found their application in various types of pheromones, agriculture. When combined with traps, luring insects can destroy a significant number of pests.*

*Keywords: physical environmental factors, environment, pheromones, highly volatile substances, releasers, primers, stimulants, pesticides, agricultural chemicals, agroecology, environmental problems, chemical pollution of the environment, research perspective, synthetic pheromones, pheromone signals, pheromone traps, pheromone monitoring of agriculture, chemosignals, lepidoptera insects, insect pests, species specificity, attractiveness, flight of insects, light, methods of pest control, methods of quantum theory, adsorption, temperature, energy, chemical formula of pheromones, atomic structure of pheromone, geometry of pheromones, electronic structure of pheromones, molecule of pheromone, conformer, limiting epoxide, unsaturated hydrocarbons, unsaturated oxygen-containing pheromones, Coulomb interaction, disparity.*

### INTRODUCTION

In foreign countries of the 20th and 21st centuries, "the problem of environmental safety has gone beyond the national and regional and has become a global problem for all mankind. Mankind really felt the threat it faces, which results in an anthropogenic impact on the environment." has brought the world to the brink of environmental disaster. Intensive human economic activity Human impact on the environment is multifaceted. The main anthropogenic factors destroying the environment are: urban growth, mining, road transport, industry and agricultural chemicals. In the deterioration of the environment, chemical exposure is in the first place. The role of chemical objects



in human life can hardly be overestimated. They are assigned one of the important places in the fight against pests, diseases and weeds of agricultural crops, but the effects of pesticides are never unambiguous. Pesticides used in agriculture are organic compounds that are toxic not only to pests, but also to humans and animals. Man uses pesticides to destroy a limited number of organisms, which make up no more than 0.5% of the total number of species inhabiting the biosphere, while pesticides, when applied, affect all living organisms. When carrying out protective measures, pesticides are always directed against the population. [2] This article examines the influence of physical environmental factors on pheromones, provides examples of studies of this kind of influence on pheromones of lepidoptera insects, as well as the prospects for the use of pheromones in agriculture. The President of the Republic of Uzbekistan Sh.M. Mirziyoyev, speaking at the meeting of the heads of state of the founders of the International Fund for Saving the Aral Sea, noted that it is necessary to start the introduction of environmentally friendly technologies, lay the foundation for the comprehensive implementation of a green economy, environmentally friendly, energy and water saving technologies. The head of state also initiated the organization of a conference next year with the support of the UN, the World Bank, the Asian Development Bank and the Global Environment Facility, at which negotiations on practical issues of creating a zone of environmental innovations and technologies in environmentally disadvantaged regions of the world will be held. This is provided for in the strategy of actions for five priority areas of development of the Republic of Uzbekistan for 2017-2021 in subparagraph 3.3 [1]. Modernization and intensive development of agriculture is understood as the implementation of investment projects for the construction of new, reconstruction and modernization of existing processing enterprises, the widespread introduction of intensive methods in agricultural production, primarily modern water and resource-saving agricultural technologies, the use of high-performance agricultural machinery; expansion of research work on the creation and introduction into production of new breeding varieties of agricultural crops resistant to diseases and pests, adapted to local soil, climatic and environmental conditions, and animal breeds with high productivity. [4]

### **THE CONTENT OF THE ARTICLE**

Insect pheromones, discovered more than fifty years ago, can now become a safe and harmless substitute for pesticides and other harmful chemicals that are currently used in the fight against harmful insects that cause huge damage to agriculture.

The question arises, what is the essence of pheromones? How do they act on insect pests? How external environmental factors affect them, and most importantly, what are the prospects for the use of pheromones in agriculture.

Pheromones are biologically active substances, products of external secretion, secreted into the environment by insects, fish, animals, communication between organisms of the same species are some kind of volatile chemosignals capable of controlling neuroendocrine behavioral reactions, developmental processes, social behavior and reproduction. Pheromones alter the behavior, physiological and emotional state, and

even the metabolism of different individuals of the same species. These substances, are means of regulation, play an important role in the communication of many species of insects, for example, ensuring the rapprochement of males and females during the breeding season, the concentration of insects on forage plants and in wintering places, or controlling the behavior and physiological processes of working individuals of social insects. Pheromones are found in animals of various taxonomic groups, from invertebrates to mammals. At present, insect pheromones are considered the most studied.

There are two main types of pheromones that differ in their effects: releasers and primers. The first type - which transmit danger signals between individuals of the same species. Usually releasers are human. Pheromones, providing chemical releasers, are able to induce an individual to take immediate action, for example, pheromones, highly volatile substances that spread through the air. The second type of the formation of special behavior and influence on other individuals, an example is the pheromones secreted by the queen bee, in order to suppress the sexual development of female bees, turning them into ordinary working bees. Primers are most commonly distributed by contact. At this time, releasers have been studied better than primers, by their example, several subtypes of pheromones can be distinguished, such as: attractants - these include sex pheromones and aggregation pheromones that stimulate the accumulation of insects; repellents - repelling pheromones; primers designed to stimulate pheromones that induce activity, for example, anxiety pheromones; deterrents that inhibit the reaction, etc. [5]

The source of pheromone in insects can be individual secretory cells scattered throughout the body or groups of them, which form a special organ of the pheromone gland. The ducts of the pheromone glands open on the surface of the body or in cavities that communicate with the external environment. Insects secrete pheromone in trace amounts: for example, a female codling moth (*Cudiapomonella*) releases only 9 nanograms of pheromone per hour. However, even this amount is enough for a male moth to smell and find a female in the crown of a tree. Insects perceive the smell of pheromones with the help of chemoreceptor sensilla - special receptors in the form of hairs, bristles or tubercles located on the antennae; their number on one antenna can reach 15 thousand. A very small amount of pheromone in the air is enough for an insect to respond. [4]

Usually, pheromones are not one substance, but a mixture of the main, predominant by weight component with small additives (minor components): they can contain more than 10 components. One substance can have several different functions. Pheromone molecules are highly volatile, quickly decompose under the action of atmospheric oxygen, moisture and light. In terms of chemical composition, insect pheromones belong to various classes of organic compounds, such as alcohols, ethers, terpenoids, steroids, aldehydes, heterocyclic compounds, and others.

Knowing the chemical composition of the insect pheromone, it is possible to synthesize it in laboratory conditions. It is these synthetic analogs of sex and aggregation pheromones that can be used to protect plants from pests. The advantage of synthetic pheromones,

which are used in micro doses, is their high species specificity and attractiveness. They are completely harmless to humans and the environment, and also act directly on the target species of insect pest.

There are two main areas of application of synthetic pheromones against harmful insects: First, monitoring. This suggests that the use of pheromones provides an opportunity to record such processes as the flight of pests, to obtain data on their numbers, or even the ability to determine the area of quarantine pests. Second, it provides an opportunity to prevent males from finding females, attract insects, and catch or destroy pest control. By saturating the air with synthetic pheromone, before they can find a natural source of pheromone. In both cases, the reproduction of pests is blocked.


However, in addition to the effect of pheromones on pests, it is necessary to consider the influence of environmental factors on the pheromones themselves. Taking into account the huge species diversity of insect pests and the complexity of the composition of pheromones, an urgent task is to develop universal methods for studying pheromone communication, which will save material, labor and time resources.

It is important to pay attention to another aspect of the possible use of pheromones - the establishment of the species composition of insects in a particular area. This is most clearly seen in the example of a scoop. Analogs of the sex pheromones of many species of scoop were synthesized. A sample of the VNIKhSZR synthesis was chosen. The observations were carried out in the cotton crop rotation of three farms in the Yangiyul district, as well as in the fields of the Scientific Research Institute of Vegetable, Melons and Gourds in the Tashkent District of the Tashkent Region. We used the pheromone-wintering scoop of two- and three-component (*Agrotissegetum*) (OTs-77 and OTs-8), exclamation scoop (*Agrotisexclamationis*) (BK-23 BK-137), C-black scoop (*Hestia c-nigrum*) (SCh-72), bindworms (*Emmeliatrabealis*) (Minus-21), cotton moths (*Helicoverpaarmigera*) (KC), meadow moths (*Mythimpainipuncta*) (MC).

Dispensers with pheromones were placed in triangular traps made of laminated paper, which were placed in the fields at the rate of 1 trap per hectare at a height of 25 cm above the plants. The pipettes were renewed every 10 days. Observations were carried out over three years in the fields of cotton, kenaf, corn, alfalfa, red pepper, tomato and pumpkin. Based on the number of males of each species caught in pheromone traps, we calculated the indicators of the relative abundance of the species [4].

## The number of the moth complex in agrocenoses of the cotton crop rotation

On the surveyed fields of cotton crop rotation (cotton, kenaf, corn, alfalfa), the complex of scoops, determined by the presence of pheromones, as differences were observed in individual years, as a rule, it is of the same type. which concerned mainly small species. Thus, in all areas the dominant species was the bindweed (*Emmeliatrabealis*), the subdominant exclamation moth (*Agrotis exclamationis*) (*Agrotissegetut*). There were no caradrina moths (*Spodopteraexigua*) and corn and winter moths (*Spodopterafrugiperda*) in the corn and cotton fields. In a cotton field, bindweed (*Emmeliatrabealis*), winter moth (*Agrotissegetum*), (*Agrotis exclamationis*), cotton moth (*Helicoverpaarmigera*), meadow moth (*Mythimnaunipuncta*), as well as gamma moth (*Autographagatta*), c.black(*Xestiac-nigrum*), ipsilon owl (*Agrotisipsilon*). The species diversity of moths in the cornfield was slightly less exclamatory: there was no cotton moth (*Helicoverpaarmigera*) and no upsilon moth (*Agrotisipsilon*).



In the alfalfa field, all types of scoops were identified, the pheromones of which were used. All kinds of scoops, whose pheromones were used during observations, (*Spodopteraexigua*) and (*Spodopterafrugiperda*) were also found in the fields of vegetable crops. In the Tashkent region, the number of scoops was in general, with the exception of caradrins of leafy corn scoops. On vegetable crops it was higher than in cotton crop rotation fields in Yangiyul region. In vegetable crops, as well as in agrocenoses of the Cotton crop rotation, the dominant species was the bindweed (*Emmeliatrabealis*), the subdominant (*Agrotis exclamationis*) was the exclamation winter (*Agrotissegetum*). So, one day after the installation of pheromone traps, 14.7 individuals were caught on red peppers, and 11 individuals were caught for tomatoes and alfalfa. At the same time, in the fields of alfalfa, tomatoes and red pepper, it was found on traps with winter pheromones of 6, 7.7 and 10.7 individuals, respectively. According to the available data, the generalized economic threshold of harmfulness is considered to be capture, on



average, per one trap per day (night) (Agrotissegetum), which corresponds to a caterpillar density of 2.6 - 4.0 individuals per 1 m<sup>2</sup>.





Diagram 1.

In the fields of vegetable crops studied by us, the number of butterflies of the winter moth (Agrotissegetum) exceeded the aforementioned EVP. With the help of pheromones from other moths, a large number of other species have been identified, which are not inferior to the winter moth (Agrotissegetum); the total number of identified moths significantly exceeded the generalized damage threshold established for only one species.



- I - winter tree(Agrotis segetum)
- II - exclamation point(Agrotis exclamationis)
- III - вьюнковаясовка(Emmelia trabealis)
- IV - луговаясовка(Mythimna unipuncta)
- V - совка-гамма(Autographa gamma)
- VI - ипсилонсовка(Agrotis ipsilon)

Chart 2.

Thus, the use of gender (sex) analogs of pheromones makes it possible to establish the specific composition of moths in the fields of various crops, as well as to identify the total number of pests in a separate field and give a signal about the need for protective measures to regulate their number. [3] The effect on pheromones, in particular on the change in their structure (destruction) from a physical point of view, is exerted by the values of the electric dipole moment ( $\mu$ , D) and wavelengths corresponding to the absorption maximum ( $A_{max}$ , nm). For oxygen-containing pheromones, the values of the electric dipole moment are in the range of 1.23-2.71 D, for unsaturated hydrocarbons, in the range from 0.29 to 0.49. Pheromones, in which partial positive and partial negative charges are located on different parts of the molecule, are mutually oriented in space so that other polar molecules are nearby - air components, for example, water molecules, whose dipole moment is 1.85 D. Coulomb interaction between polar fragments of molecules can create favorable conditions for their mutual orientation, the occurrence of a chemical reaction and, as a consequence, for the destruction of pheromone. Modeling of photo-excitation processes has shown that the presence, number and mutual arrangement of multiple bonds in a molecule affects the absorption wavelengths. For pheromones that do not contain multiple bonds, the absorption maximum lies in the range 136-144 nm, for unsaturated hydrocarbons and unsaturated oxygen-containing pheromones, in the range 157-204 nm, for oxygen-containing pheromones with conjugated double bonds, in the range 226-230 nm. The intensity of the pheromone signal will decrease when exposed to light radiation, since the excited state is unstable, the molecule goes into the ground state, losing energy due to radiation, vibrations or collision with other molecules. changes in the electronic and atomic structure Exposure to electromagnetic radiation leads to pheromones. Most pheromones of lepidoptera insects contain up to three double bonds in the structure. Analysis of the electronic structure of pheromone molecules shows that, regardless of the presence and type of the oxygen-containing functional group, the redistribution of the electron density upon absorption of light occurs in the region of the location of double bonds and corresponds to the transition of an electron from n-bonding to n\* -brobing orbitals. The change in bond lengths occurs only in the region arrangement of multiple bonds and does not affect oxygen-containing functional groups. Such a change in the bond length is unlikely to lead to the destruction of the initial atomic structure of molecules only under the influence of electromagnetic radiation, but will increase their reactivity and promote chemical reactions when interacting with air components. For bicyclic pheromones, upon transition to an excited state, an increase in the length of one of the bonds included in both cycles is observed, which can lead to the opening of one of the cycles of the molecule and the destruction of the pheromone molecule. An assessment of the resistance of pheromones to thermal exposure using the results of ab initio calculations showed that thermal exposure in noncyclic and monocyclic molecules is most likely to change the lengths of single bonds, while light radiation primarily affects multiple bonds, leading to an increase in their lengths. For bicyclic unsaturated pheromones, exposure to light radiation and thermal effects lead to similar structural changes - an increase in the length of a single bond included in both

cycles, which can lead to its rupture and the opening of one of the cycles in the structure of the molecule. The factors that can deactivate pheromone molecules can be adsorption on the surface of plants, exposure to temperature, light and chemical interaction with substances-components of the air. Water molecules that are present in the air can interact with polar pheromone molecules, which will lead to a decrease in their concentration. Exposure to light radiation of the ultraviolet part of the spectrum transforms pheromone molecules into an excited state, while the bond lengths and bond angles in the molecules change. Connecting the considered process of excitation of a molecule with communication of insects, it can be assumed that a pheromone molecule in an excited state may not be captured by insect sensors, since the efficiency of binding of a pheromone inside a receptor depends on its atomic structure, that is, the intensity of the pheromone signal will decrease. Pheromone molecules must have certain physical characteristics in order to remain in the air flow for some time, sufficient for signal propagation, but they must not accumulate in the territory to carry up-to-date information on the position of a variety of insect pests, a multicomponent composition of pheromones, complex multistage processes of obtaining pheromones from alternative non-experimental methods for studying the properties of pheromones and mechanisms of chemical communication. We can examine the properties of pheromone molecules of insect pests by examining them using the methods of the quantum theory of condensed matter. The aim of the study was to theoretically study the physical properties of disordered organic systems, namely the atomic and electronic structure of pheromone molecules in the ground and excited states, to identify their photophysical characteristics, and to assess their stability under light and thermal exposure. The application of the methods of quantum theory allows one to obtain the basic physical characteristics of the pheromone signal source. A huge species in laboratory conditions lead to the need to develop insect molecules and evaluate their resistance to such external factors as thermal and light exposure without the use of expensive experimental techniques. This approach is universal for different types of pheromone molecules of various insect species, which is very important, given the multicomponent composition of pheromones and the species diversity of insect pests. Establishing the relationship between the search behavior of insects and the physical characteristics of pheromone molecules can be used to obtain information about the search behavior of insects, based on data on the chemical composition of pheromone. The data on the resistance of pheromone molecules to the action of environmental factors, obtained using *ab initio* methods of quantum theory, can be used to modernize and increase the efficiency of methods for controlling the number of insects based on the use of pheromone drugs.

#### RESEARCH METHODS

Theoretical modeling and calculation of the physical characteristics of pheromone molecules were carried out within the framework of the density functional theory using the B3LYP functional, using the 6-31G \*\* and cc-pVDZ basic packages, implemented in the GAMESS-US program. The absorption spectra were calculated and the molecules were optimized in the excited state by the TimeDerecent method. The assessment of the



Tipferomone	$\mu, D$	$\lambda_{max, HM}$
Pheromone, not containing multiple bonds	1,76-2,04	136-144
Unsaturated hydrocarbons	0,29-0,49	179
Unsaturated oxygenated pheromones	1,23-2,65	168-204
Unsaturated oxygenated pheromones with conjugated double bonds	1,50-2,71	224-227

Unsaturated oxygenated pheromones with  
conjugated double bonds 1,50-2,71 224-227

Table 1. Calculated values of the electric dipole moment ( $\mu$ , D) and wavelengths corresponding to the absorption maximum ( $\lambda_{max}$ , nm) for pheromones of lepidopteran insects.

For oxygen-containing pheromones, the values of the electric dipole moment are in the range of 1.23-2.71 D, for unsaturated hydrocarbons, in the range from 0.29 to 0.49. Pheromones, in which partial positive and partial negative charges are located on different parts of the molecule, are mutually oriented in space so that other polar molecules - air components, for example, water molecules, whose dipole moment is 1.85 D. Coulomb interaction between polar fragments of molecules can create favorable conditions for their mutual orientation, the occurrence of a chemical reaction and, as a consequence, for the destruction of pheromone. Considering the high polarity of molecules as a factor promoting the chemical interaction of pheromones with polar components of the medium, it can be concluded that for oxygen-containing pheromones the probability of destruction as a result of chemical reactions will be higher than for pheromones-hydrocarbons, whose dipole moment is significantly less than 1 D, in all possible conformations.

All calculated pheromones of Lepidoptera absorb radiation corresponding to the ultraviolet part of the spectrum, in the range from 130 to 230 nm. Changing the geometry for some pheromones leads to a shift in the absorption maximum, but by no more than 8 nm, that is, no more than 6%. Analysis of the spectral characteristics showed that the maximum absorption of pheromones depends primarily on the presence and mutual arrangement of multiple bonds and practically does not depend on the type of oxygen-containing functional group. The greatest energy is required for the excitation of the limiting epoxide, the minimum for pheromones with conjugated double bonds.

As noted earlier, a change in the conformation of the molecule does not lead to a significant shift in the absorption maximum.




Figure 3 - Atomic and electronic structure of pheromones in the ground and excited state: carbon atoms are marked in blue, hydrogen atoms - in gray, oxygen atoms - in red.

Exposure to electromagnetic radiation leads to a change in the electronic and atomic structure of pheromones. For disparlyur, a pheromone of the gypsy moth (*Lymantria dispar*), which does not have multiple bonds in the structure, the change in the electron density upon absorption of radiation occurs in the region of the epoxy ring for all conformers. Calculation of the geometry of the disparlyur in an excited state shows similar structural changes for all conformations: the bond angles change in the epoxy ring, one of the C-O bonds increases on average to 0.9 Å, which can further lead to its rupture and, as a consequence, to the opening of the cycle and the destruction of the pheromone. Most pheromones of lepidoptera insects belong to unsaturated compounds containing up to three double bonds in their structure. An analysis of the electronic structure of pheromone molecules shows that, regardless of the presence and type of oxygen-containing functional group, the redistribution of the electron density upon absorption of light occurs in the region of the location of double bonds and corresponds to the transition of an electron from n-bonding to n\* -bonding orbitals (Figure 3). The bond lengths are indicated in angstroms (Å).

Figure 3 shows the values of the lengths of chemical bonds in molecules of unsaturated alcohol and unsaturated epoxide in the ground and in the excited state. The change in bond lengths occurs only in the area of multiple bonds and does not affect oxygen-containing functional groups. Similar structural changes occur in all unsaturated pheromones of Lepidoptera. An increase in the lengths of double bonds occurs on average by 0.1 Å. Such a change in the bond length is unlikely to lead to the destruction of the initial atomic structure of molecules only under the influence of electromagnetic radiation, but will increase their reactivity and promote chemical reactions when interacting with air components. The interaction of the pheromone with the protein of the insect's olfactory receptor occurs according to the "key-lock" principle, that is, in

addition to the chemical composition of the pheromone, its geometric correspondence with the protein plays an important role. Therefore, the course of chemical reactions or a significant change in the initial geometry of the molecule will lead to the deactivation of the pheromone as a carrier of information.

The calculated characteristics of lepidoptera pheromones were compared with the data on the search activity of insects.

For example, the pheromone of the gypsy moth (*Lymantria dispar*) consists of one component, *disparlura*, and the pheromone of the pine silkworm (*Dendrolimus pini*) contains four components. One can see clear differences in the intensity and wavelengths of absorption of pheromones of these two types. It is known that the gypsy moth (*Lymantria dispar*), whose pheromone absorbs at 130 nm with a very low intensity, is characterized by searching behavior throughout the day, while the pine silkworm (*Dendrolimus pini*) is characterized by activity in the evening and night hours. in the absence of solar radiation. For pheromones of the Siberian silkworm (*Dendrolimus sibiricus*) and moths (*Geometridae*) of moths, the absorption spectra lie in the same range as for the pine silkworm (*Dendrolimus pini*) and have similar values of the intensity of electronic transitions. The traced relationship between the spectral characteristics of molecules and data on the flight time suggests that the physical properties of pheromones to some extent determine the timing of the search activity of insects. [6]

#### CONCLUSION

The efficiency of information transfer using pheromone molecules is determined by a number of factors, for example, such as the resistance of pheromones to the effects of the external environment on them, that is, to their physicochemical characteristics. The purpose of pheromones and the principle of their action is based on their preservation of their composition and structure, for a certain time, which should be sufficient for spreading in the air, and reaching individuals that must receive a chemical signal. And the use of highly resistant molecules as pheromones can lead to clogging of the information channel and disorientation of individuals receiving signals.

## Used literature

1. Decree of the President of the Republic of Uzbekistan "On the Strategy of Action for Five Priority Directions for the Development of the Republic of Uzbekistan in 2017-2021" UP No. 4947 of February 7, 2017 // Teacher of Uzbekistan. - T. - No. 6 (2453). - February 10th, p. 2.

2. Khonkhodzhaeva Nodira Bakhtiyarovna. Ecological features of the use of monitoring in agriculture. // European Journal of Research and Reflection in Educational Science Vol. No. 8. 2020

3. Khonkhodzhaeva N.B. (Republic of Uzbekistan) PHEROMONS AND THEIR BIOLOGICAL SIGNIFICANCE

4. Details.uz: NewsAgency. August 13, 2019 (Date of access 08.13.2019).

5. Dmitrieva T.M., Kozlov Yu.P., Pheromones AND THEIR ROLE IN

