



BRITISH

MEDICAL JOURNAL



British Medical Journal Volume-2, No 1

10.5281/zenodo.6368043

British Medical Journal

Volume 2, No 1., 2022

Internet address: <http://ejournals.id/index.php/bmj>

E-mail: info@ejournals.id

Published by British Medical Journal

Issued Bimonthly

3 knoll drive. London. N14 5LU United Kingdom

+44 7542 987055

Chief Editor

Dr. Fiona Egea

Requirements for the authors.

The manuscript authors must provide reliable results of the work done, as well as an objective judgment on the significance of the study. The data underlying the work should be presented accurately, without errors. The work should contain enough details and bibliographic references for possible reproduction. False or knowingly erroneous statements are perceived as unethical behavior and unacceptable.

Authors should make sure that the original work is submitted and, if other authors' works or claims are used, provide appropriate bibliographic references or citations. Plagiarism can exist in many forms - from representing someone else's work as copyright to copying or paraphrasing significant parts of another's work without attribution, as well as claiming one's rights to the results of another's research. Plagiarism in all forms constitutes unethical acts and is unacceptable.

Responsibility for plagiarism is entirely on the shoulders of the authors.

Significant errors in published works. If the author detects significant errors or inaccuracies in the publication, the author must inform the editor of the journal or the publisher about this and interact with them in order to remove the publication as soon as possible or correct errors. If the editor or publisher has received information from a third party that the publication contains significant errors, the author must withdraw the work or correct the errors as soon as possible.

OPEN ACCESS

Copyright © 2022 by British Medical Journal

British Medical Journal Volume-2, No 1

ADVANTAGES OF ULTRASONOGRAPHY IN INFLAMMATORY AND DEGENERATIVE DISEASES OF THE HIP JOINT

Madumarova Zarnigor Shukhrat kizi

Andijan State Medical Institute, Faculty of Advanced Training and Retraining of Physicians, senior teacher at the Department of Medical Radiology and Clinical Laboratory Diagnostics,
Andijan city, Uzbekistan
zarnigor-nazarova@ yandex.ru

Nazarova Gulchexra Usmonovna

PhD Andijan State Medical Institute, Faculty of Advanced Training and Retraining of Physicians, Head of the Department of Family Medicine, Andijan city, Uzbekistan

mjpand@mail.ru

Khojibekov Marat Khudoykulovich,

Doctor of Medical Sciences, Professor, Tashkent Medical Academy

Abstract: This article is devoted to the application of ultrasonography in the diagnosis of inflammatory and degenerative diseases of the hip joint. The article discusses the relevance of the use of ultrasound in the disease of osteoarthritis (OA), rheumatoid arthritis and aseptic necrosis of the femoral head, as well as the differential assessment of inflammatory and degenerative processes based on ultrasound Doppler sonography. For a long time, a simple X-ray was considered the reference technique of osteoarthritis (OA). Recently, ultrasonography is an innovative method for visualizing this disease. The use of ultrasound highlights the various anatomical structures in great detail and detects intra- and extra-articular changes.

Keywords: Coxoarthrosis, coxitis, rheumatoid arthritis, osyearthritis aseptic necrosis, ultrasonography, hip joints, diagnostics.

Introduction

Hip joint disease is one of the urgent problems of modern orthopedics, as it contributes to the rapid development of severe anatomical and functional inferiority of the musculoskeletal system, in particular in the elderly. According to WHO, more than ten percent of the world's population suffers from joint diseases. In terms of the frequency of lesions, the hip joint ranks first (42.7%), the second is the knee (33.3%), the third is the shoulder (10.8%), and the remaining joints account for 13.2% [1].

The differential diagnosis of lesions of the hip joint is difficult due to its deep occurrence and the presence of large muscle masses around the joint. In this regard, it is not possible to assess the external changes in the joint, it is very difficult to perform its puncture and arthroscopy. The differential diagnosis of coxarthrosis and coxitis at an early stage is especially difficult [2]. In this regard, the issue of differential diagnosis of diseases of the hip joint of dystrophic and inflammatory origin is relevant.

The ultrasound method, being non-invasive, reproducible and relatively affordable, can be widely used to assess both the accumulation of fluid inside the joint, in the joint bags, and to assess the thickness of the synovial membrane and erosive changes [3,4]. Modern ultrasound equipment makes it possible to evaluate both superficial periarticular and intra-articular tissues that make up the joint.

Purpose of the study: Improving the Efficiency of Radiation Diagnosis of Inflammatory and Degenerative Diseases of the Hip Joint Based on the Ultrasound Method.

Materials and methods of research.

The total number of patients examined was 138, of which 15 (13.9%) were men and 93 were women (86.1%). The patients were divided into groups and subgroups. Group I included patients with dystrophic diseases (coxarthrosis stage I-III), a total of 82 people. Group II included 10 patients with coxarthrosis formed as a result of aseptic necrosis of the femoral head. Of these, men 5 (50%) people, women 5 (50%) people. Group III with inflammatory diseases included patients with rheumatoid arthritis, in the amount of 12 people. Of these, 4 men (33.3%) people, women 8 (66.7%) people.

All patients in the period from 2018 - 2020. were treated by an orthopedist in the ASMI clinic, and an ultrasound examination was carried out in the radiology department of the ASMI clinic in Andijan.

Ultrasound examination was carried out on digital multifunctional scanners "Sonoscape S-22" (China) and "Mindray DC3" (China). For the most clear visualization, a study was used with convex and linear sensors, in the frequency range of 7-12 MHz.

Comprehensive ultrasound examination included survey scanning of the hip joint in gray scale mode, examination of the shape and contour of the femoral head, determination of the presence of effusion, measurement of parameters of the joint capsule, thickness of the hypoechoic layer of hyaline cartilage, examination of the state of muscles, ligaments and bones adjacent to the joint.

When evaluating the structures of the hip joint according to the standard method from the anterior approach, the bone landmarks were the upper edge of the acetabulum and the semicircle of the femoral head, where hypoechoic hyaline cartilage, the articular capsule of the hip joint were visualized (Fig. 1).




Fig. 1. Sonogram of the structural elements of the hip joint is normal. The arrows in the diagram show the measurement of the thickness of the hypoechoic layer of hyaline cartilage. 2 - fibrous capsule.

In patients of all groups, the regional blood flow in the lateral circumflex arteries of the thigh was assessed. The assessment of blood flow was carried out using color Doppler mapping and pulsed wave spectral Doppler, while assessing: peak blood flow velocity (PSV), end-diastolic blood flow velocity (DPV), resistance index (IR).

Research results

When analyzing the revealed sonographic and Dopplerographic diagnostic criteria for the stages of the pathological process, in patients of group I, it was revealed that the thickness of the hypochoic layer of hyaline cartilage decreases from normal values of 2.4 mm to 1.3 mm as the stages of the disease increase in patients of subgroup IA, to 0.9 mm in subgroup IB and thins to 0.8 mm in subgroup I C. At stage III of coxarthrosis, the thickness of the hypochoic layer of hyaline cartilage is 0.85 mm, and with coxarthrosis formed as a result of aseptic necrosis of the femoral head, cartilage in the necrosis zone is not visualized. In rheumatoid arthritis, there was an increase in the thickness of the fibrous capsule, up to 3.4 mm, in contrast to 2.23 mm with unchanged joints, and an important sign was the identification of normal sizes of hyaline cartilage. At the same time, the sizes of osteophytes increased as the stage of coxarthrosis increased from 4.2 mm to 10.1 mm. Thus, in stage III coxarthrosis, the size of osteophytes was 8.3 mm, and in coxarthrosis formed as a result of aseptic necrosis of the femoral head, it increased to 15.0 mm.

All patients with stages I-III of coxarthrosis revealed significant differences in the shape of the head of the hip joint, so if at stage I of the disease almost all (96%) of the examined patients had a spherical shape of the head, then at stage II of the disease only in 58% of cases a spherical shape was detected, and in 42% of cases - a moderately flattened shape of the head. At stage III, this sign increased and in 96% of cases it was found to be significantly flattened (Fig. 2).



Fig.2. Sonogram of the hip joint in coxarthrosis (significantly flattened shape of the head).

When studying the joint cavity in patients with stage I coxarthrosis, effusion was practically absent (1% of cases), at stage II this sign was already detected in 10% of cases, and in patients with stage III coxarthrosis, effusion was visualized in 37% of cases. That is, as the disease progresses, the likelihood of effusion in the joint cavity increases.

When applying the method of ultrasonic dopplerography, a progressive decrease in the speed indicators of blood flow in the lateral circumflex arteries of the thigh was revealed as the stage of coxarthrosis increases, so the PVR decreases from

25.2 cm/s in the first stage of coxarthrosis to 16.6 cm/s in the third stage; CDS from 4.6 cm/s at the first stage to 2.1 cm/s at the III stage of coxarthrosis. In addition, if the PSS at stage II is 22.4 cm/s, and with coxarthrosis formed as a result of aseptic necrosis of the femoral head, it significantly decreases to an average of 17.8 cm/s. In rheumatoid arthritis, there was an increase in PSS of blood flow, on average, up to 28.0 cm/s, as well as an increase in DPV of blood flow, up to 8.1 cm/s. and a decrease in IR to 0.65, when comparing the values of groups with unchanged joints and stage II of coxarthrosis at 0.78 and 0.77, respectively.

Discussion

The thickness of the hyaline cartilage and fibrous capsule, the presence or absence of intra-articular effusion, the shape of the femoral head, the size of osteophytes (if any) are the main structural parameters on which the differential diagnosis of dystrophic and inflammatory diseases of the hip joint is based. In case of an inflammatory lesion of the hip joint, the leading sonographic criteria are: an increase in PVR to numbers - 32.0 cm/s, a CDP to - 10.3 cm/s, and a decrease in IR to numbers - 0.65, combined with an increase in the thickness of the fibrous capsules while maintaining the normal thickness of the hyaline cartilage. And the signs of the presence of coxarthrosis are: a progressive decrease in the rate of blood flow in the lateral circumflex arteries of the thigh and progressive thinning of the hyaline cartilage, the presence of marginal osteophytes, a violation of the shape and contour of the femoral head. Coxarthrosis, formed as a result of aseptic necrosis of the femoral head, is characterized by a significant decrease in the rate of blood flow in the lateral circumflex arteries of the thigh (PRS up to digits - 17.8 cm/s, CDS up to digits - 3.7 cm/s) in combination with early and more complete thinning of the hyaline cartilage, more significant sizes of osteophytes, significant deformity of the femoral head, effusion in the joint cavity.

Conclusions

A number of undoubted advantages - non-invasiveness (unlike arthroscopy), accessibility, simplicity, cost-effectiveness (compared to CT and MRI) - provided the method of hip joint ultrasound with a priority among other instrumental methods for examining joints and soft tissues [5]. Ultrasound is highly informative in reflecting small details of the surface of bones, ligamentous tendon apparatus, and also allows you to identify and control inflammatory changes in tissues. The advantage of ultrasound over the X-ray method is safety, polypositionality, as well as the possibility of dynamic observation.

The most important features of a comprehensive ultrasound examination of the hip joints are: determination of the state of the joint capsule, the presence of effusion in the cavity of the hip joint, assessment of the sphericity and contour of the head, determination of the state of the hyaline cartilage and periarticular tissues, and using the Doppler method, the blood flow in the lateral circumflex arteries of the thigh is assessed, with the calculation of blood flow PSV, blood flow DPV and IR, which is quite important in the differentiation of inflammatory and degenerative processes.

References

1. Zhang, W. EULAR recommendations for knee and hip osteoarthritis: a critique of the methodology / W. Zhang, M. Doherty // *Br. J. Sports Med.* - 2006. -Vol. 40.-p.

2. Kovalenko, V.N. Osteoporosis. Practical guidance /N. Kovalenko, O.P. Bortkevich. - K.: Morion, 2003. - 448 p.
3. Khabirov, R.A. Ultrasound differential diagnosis of degenerative and inflammatory joint lesions / R.A. Khabirov, M.G. Tukhbatulin, E.R. Volkova // Nauch. - pract. rheumatology. - 2001. - No. 3. -p. 125.
4. Yaremenko, O.B. Modern algorithm for diagnosing rheumatoid arthritis / O.B. Yaremenko // Clinical Immunology. Allergology. Infectology. - Kiev, 2006. - No. 2. - p. 54-58.
5. Saltykova, V.G. Possibilities of ultrasound diagnostics in injuries and diseases of large joints / V.G. Saltykova, A.K. Morozov // Proceedings of the All-Russian Congress of Radiation Diagnostics. - M., 2007.-p. 316-317.
6. Divakov M.G., Boloboshko K.B. Early ultrasound diagnosis of osteochondropathy of the femoral head // News of Radiation Diagnostics. 1999; 2:12-13.