



BRITISH

MEDICAL JOURNAL



British Medical Journal Volume-2, No 4
10.5281/zenodo.7109681
British Medical Journal
Volume 2, No 1., 2022
Internet address: <http://ejournals.id/index.php/bmj>
E-mail: info@ejournals.id
Published by British Medical Journal
Issued Bimonthly
3 knoll drive. London. N14 5LU United Kingdom
+44 7542 987055
Chief Editor
Dr. Fiona Egea

Requirements for the authors.

The manuscript authors must provide reliable results of the work done, as well as an objective judgment on the significance of the study. The data underlying the work should be presented accurately, without errors. The work should contain enough details and bibliographic references for possible reproduction. False or knowingly erroneous statements are perceived as unethical behavior and unacceptable.

Authors should make sure that the original work is submitted and, if other authors' works or claims are used, provide appropriate bibliographic references or citations. Plagiarism can exist in many forms - from representing someone else's work as copyright to copying or paraphrasing significant parts of another's work without attribution, as well as claiming one's rights to the results of another's research. Plagiarism in all forms constitutes unethical acts and is unacceptable. Responsibility for plagiarism is entirely on the shoulders of the authors.

Significant errors in published works. If the author detects significant errors or inaccuracies in the publication, the author must inform the editor of the journal or the publisher about this and interact with them in order to remove the publication as soon as possible or correct errors. If the editor or publisher has received information from a third party that the publication contains significant errors, the author must withdraw the work or correct the errors as soon as possible.

OPEN ACCESS

Copyright © 2022 by British Medical Journal
British Medical Journal Volume-2, No 4

ELSEVIER



SSRN
Tomorrow's Research Today

Universal
Impact Factor



Morphological changes in neuroimmunoendocrine cells off the small intestine in the early postnatal period of life

Makhmudova Ziyoda Takhirovna, Talipov Rustam.

Center for the Development of Professional Qualifications of Medical Workers, The Republic of Uzbekistan, Tashkent.

Tashkent state dental institute, The Republic of Uzbekistan, Tashkent.

E-mail: makhmudova-zieoda@mail.ru

Abstract: This article is devoted to studying the features of the absorption process and the regulation of homeostasis in the early postnatal period. The material of the study was outbred white rats, which were on different types of feeding. The small intestine with normal microflora affects the adequate adaptation of functional systems of various levels of organization, metabolism, normal structure and function of all organs and systems, the body as a whole.

Keywords: Homeostasis, adaptation, microbiocenosis of the small intestine, neuroimmunoendocrine system.

It is known that natural, mixed or artificial, definitive nutrition involves the intake of nutrients and a huge number of various microorganisms from the external environment into the intestine. As a result, during the formation of a normal intestinal microbiocenosis, a genetically determined harmonious development of the individual is ensured. The formation of intestinal microbiocenosis is significantly influenced by childbirth (natural or caesarean section), microbiocenosis of the birth canal, ecology, drugs, especially antibiotics, nosocomial infection, type of feeding, etc. [2,3,10.].

However, to date, existing ideas about the role and importance of the intestinal microflora in the formation of the small intestine, its neuroimmune - endocrine formations, the functional system of the external environment-microbiocenosis intestines - the internal environment of the microorganism, adaptation and regulation of homeostasis in the process of digestion and absorption are fragmented and contradictory.

Material and research methods. Rats are white outbred at the age of 1.3, 7 and 14 days after birth, which were breastfed. Slaughter of animals and taking a piece of the initial section of the jejunum was carried out in accordance with the International Convention for the Protection of Animals used for Scientific Purposes (2003). After appropriate fixation and wiring, obtaining ultrathin sections, the material was viewed in an IEM - 100 S electron microscope. In the dynamics of age, the activity of hydrolytic enzymes in the homogenate of the mucous membrane of the jejunum was determined by conventional biochemical methods.

Research results.

In newborn rats, as well as children [6,9,11], before feeding, the mucous membrane of the small intestine is not separated from the submucosa, forms villi of various generations and short rare crypts between them. Formed finger-like villi are expelled by highly prismatic epithelium, have a homogeneous cytoplasm and a wide, up to 1.0 μm brush border on the apical surface. Goblet cells between them are

single, have a characteristic ultrastructure and secretory granules of moderate density in the supranuclear region. Endocrine and neuroreceptor cells are rarely detected, they are at the stage of differentiation. Lymphocytes or other leukocytes are not detected between villi or crypt enterocytes. In the lamina propria under the epithelium, single small groups of cells are detected, consisting of clusters of lymphoblasts.

In newborn rats (1-3 days), which are in the natural conditions of the vivarium, due to the minimal development and differentiation of cells of the fundic glands of the stomach, pancreatic acini, low hydrolytic-transport function of columnar epithelial cells of the villi of the small intestine [9], digestion is autolytic and symbiotic, carried out in the intestinal cavity. Insufficiency of mucus production by goblet cells and developing Brunner's glands in the duodenum does not allow for practical implementation of membrane and membrane digestion in newborns and rats [9]. Absorption from the lumen of the jejunum into the cytoplasm of villi enterocytes is carried out heterochronously, by receptor-mediated endocytosis. This is a perfect mechanism for the adaptation of mammals to natural feeding with breast milk, which is carried out 0.5-1.0 hours after birth, naturally due to the termination of amniotic and placental nutrition in the same receptor-mediated way.

After feeding rat pups, regulatory, protective and other biologically active ingredients in the composition of breast milk already after 3-5 minutes enter the composition of the intestinal chyme and begin to interact with the stalk-like border enterocytes of the villi, plasmolemm receptors between the bases of the microvilli, forming tubulo-vesicular formations. As a result, using the most perfect way to maintain homeostasis of the internal environment of the newborn receptor-mediated endocytosis in a matter of seconds, the binding of plastic, protective and biologically active ingredients contained in the composition of breast milk occurs. Transport of plastic, protective and biologically active ingredients from the lumen of the organ to the cytoplasm, to the supranuclear zone, to the structures of the Golgi complex.

In man, the transition from the sterile conditions of his symbiotic development in the mother's womb to the anti-natal ones occurs against the background of continuing close relationships with the mother. Within 1-2 years, its adaptation consists in the formation of neuroimmunoendocrine and other functional systems [1,8] in close relationship with ecology, the properties of numerous microorganisms, dynamic symbiotic relationships with the external environment, a characteristic type of nutrition, digestion, trophology, which have been evolutionarily fixed in the form of harmonious integration of normal microbiocenosis of the intestine and internal environments, structural and functional adaptation of neuroimmunoendocrine and other body systems, regulation of homeostasis of the internal environment.

On the 1st-3rd day after birth, in the lamina propria of the mucous membrane of the small intestine, mainly mesenchymal and rarely monocyte-like cells, fibroblasts are detected. Blood and lymphatic capillaries are at the stage of formation and growth. Nerve elements, accumulations of lymphoblasts and reticular cells are rarely detected. In the caudal part of the duodenum and ileum, under the epithelium,

round or oval formations are detected, consisting of diffusely located lymphoblasts and reticular cells. Among them, mitotically dividing cells are often noted.

In 7 day old rats, along the mucous membrane of the small intestine with normal microflora, the proportion of formed villi and growing crypts, the number and density of endocrine cells in them significantly increase. At the same time, the degree of infiltration of the epithelial layer of villi and crypts by lymphocytes increases. In the lamina propria, the density of nerve fibers, differentiated leukocytes and connective tissue cells, and blood capillaries increases. As the submucosal and muscular layers thicken, the nerve ganglia become larger, the density and degree of differentiation of various nerve cells increase in them. In progressively increasing accumulations of lymphoblasts and reticular cells, monocytes, an emerging network of blood and lymphatic capillaries are detected.

7 days after the birth of rats with normal microflora, both the number of lymphoid formations along the organ and their volume increase. Individual follicles, or nodules, are indistinguishable. With a diffuse arrangement of lymphoblasts and reticular cells in each of the clusters of lymphoid tissue, the absolute number of cells in them increased on average 2 times. Among them, macrophages are found first, which are moderately active and contain polymorphic lysosomes.

2 weeks after the birth of rats along the small intestine with normal microflora in the mucous membrane, the proximo-distal gradient of the linear parameters of the villi and crypts becomes more distinct, the frequency of their neoplasms significantly decreases. In the thickness of the wall of the small intestine, the number of Peyer's patches reaches 10.8 ± 1.6 . In each of the accumulations of lymphoid tissue, an intensive increase in the number and density of cells leads to their bulging into the intestinal lumen, pushing the villi and crypts to the periphery. Their luminal surface is lined with a single-layer prismatic epithelium, where there are border, M and neuroepithelial cells, single goblet cells. The epithelium is infiltrated with lymphocytes at different levels. In the lamina propria and submucosa of the small intestine, in the accumulations of lymphoid tissue of the proximal part of the small intestine, germinative and follicular zones are more often seen in the distal part. Other structural and functional zones do not differ.

Structural and functional formation of the intestine occurs between 3-4 weeks after the birth of rats, when they switch to definitive nutrition. In them, as in sexually mature animals 3-4 months old with normal intestinal microbiocenosis, the mucous membrane of the small intestine, consisting of the epithelium of the connective tissue and muscular plates, has a characteristic relief due to the presence of folds, crypts and villi. The structural and functional unit of the mucous membrane of the small intestine is the crypt-villus system. It establishes certain dynamic relationships between proliferating, functioning and extruding epithelial cells. Proliferation of epitheliocytes is carried out in the lower half of the crypts. For border and goblet cells, the life cycle is on average 72 hours. The functioning of the border and goblet cells throughout the villi lasts 24-48 hours. The extrusion process can be observed at any level of the villi due to their heterochronous functioning.

Based on the study of the mucous membrane of the small intestine, the formation after the birth of symbiotic relationships between macro- and microorganisms, the regular introduction of nutrients and biologically active substrates into the body for the purpose of the harmonious development of the individual, adaptation and homeostasis of the internal environment of the normal structure and function of internal organs and systems naturally formed in the evolution of the functional system internal macroorganism environment - intestinal microbiocenosis-external environment. Feedback of the peripheral (in the small intestine) and central neuroimmune systems (hypothalamic-pituitary system), symbiotic relationships between membrane dominant and associations of cavitory microsymbionts, optimal conjugation of symbiotic cavitory and membrane, sterile membrane digestion associated with absorption into enterocytes villi provide the most important thing - homeostasis of the internal environment of the microorganism, adequate adaptation of functional systems of various levels of organization, metabolism, normal structure and function of all organs and systems, the body as a whole.

References:

1. Zufarov K.A., Yuldashev A.Yu. Pancreas., // Guide to histology. - St. Petersburg, 2001.- v.2.- S.115-141.
2. Kovalchuk L.V., Gankovskaya L.V. Lack of leukocyte prime in newborns // Immunology - 2000.-№ 3-p.12-15.
3. Mozheiko L.A. Endocrine-exocrine relationship of the pancreas: background. Journal of Grodno State Medical University 2007. P 7-11.
4. Parfenov A. I. Enterology . M.: Triada, 2002.-702 p.
5. Panegin B.V., Karsakova M.I. Macrophages: properties and function. // Immunology - 2009. - No. 3 - P. 241-249.
6. Rylova N.V. Diagnosis of diseases of the pancreas in children. Kazan. Practical Medicine - 2010
7. Sudakov KV Physiology of functional systems of the body. M.: Medicine, 2005.-304 p.
8. Terentiev A.A., Gurina A.E., Mikaelyan N.P. The state of insulin receptors in case of damage to the pancreas under experimental conditions. Moscow. Health and Education . 2014
9. Khavkin A.I. Microflora of the digestive tract. M.: Social Protection Fund, 2006.-416 p.
10. Khaitov R. M. Physiology of the immune system . M .: VINITI, 2005-448s.
11. Yuldashev A.Yu., Rakhmanov R.R., Nishanova A.A. and other. Mechanism of regulation of homeostasis in the absorption of protein from the small intestine into the blood // Medical Journal of Uzbekistan - 2009, - No. 5. - P. 79-87.
12. Yuldashev A.Yu., Kaharova Z.A., Yuldashev M.A. et al. Functional morphology of the immune system of the small intestine mucosa. Tashkent: Yangi asr avlodi . – 2008.- 48 p.